MISSOURI BOTANICAL GARDEN

PETER H. RAVEN LIBRARY

GEORGE ENGELMANN BOTANICAL NOTEBOOKS



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

7508

1-14

59 Saxifrageae

Ribesiaceae s. Grossulareae

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
Missouri Botanical Garden

Ribes leptanthum Gray

Fendl

June 1862

× 4



tube hairy

750

tube inside naked, slightly infundibuliform, much longer than ovary

lobes longer than tube, spreading

petals rather long, more than half as long as lobes

stamens same length – petals

style bifid at apex only

smooth

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

Ribes ~~[illegible]~~ hirtellum
just ripening
pulp still coheering

June 12th 1862



this part of
arillus was
imbedded in the
furrow a
arillus of
very different shape
in different seeds

751

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

JONES & LEITCH,
DRUGGISTS,
Corner Fifth and Market Streets, St. Louis, Mo.

leptanthum

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

Ribes rubrum

June 1862

ripe seed

x5

Epidermis raised

752





sect.

funicle covered by epiderm

At full maturity the cells of the epidermis become empty, the juice fills the cavity of ovary, and the former forms an empty bag. (See Saxifrag.

x60



surface of epidermis

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI BOTANICAL GARDEN

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST.
Marble Building Cor. Fourth & Olive Sts. St. Louis, Mo.

Ribes Grossularia

May 1862

753

nearly half full grown, but unripe.

x5



arillus
and funiculus

funiculi and
placentae
agglutinated
forming 2 large
white fleshy
masses

full grown
unripe

x5



x60



x5



section. [quite correct?]



over

x60



cellular tissue of Arillus
on different
planes

x5



x60



perpendicular
sect. of cells.

ripe seed, epidermis almost
dissolving, arillus very large
also dissolving into pulp

Arillus in some varieties much larger yet, enveloping greater part of seeds

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

M. W. ALEXANDER,
SOUTH-EAST CORNER OF FOURTH AND MARKET STREETS,
ST. LOUIS.

in the original section the fasciculus was ~~seen~~ thought to be seen exterior to epidermis, but it is very evidently (see also sect of Ribes rubrum) enclosed by the cells of the epidermis, though they are here much less developed — it becomes detached by the dissolving of the epidermis cells.

Ribes rubrum

May 1862 & June

754

full grown but unripe seed

×5

sc



somewhat cordate arillus

testa already transparent, thick.

×60



surface of unripe seed



epidermis, easily separated very thin and transparent.



cells after the separation of the epidermis

×60

over



×415





lobed arillus, detached

×5



Mass of seeds from below

(Raphe usually on lower side, but irregular.

Arillus cupshaped, lobed consisting of mass of (Placenta &) funiculus attached to and grown from

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
Missouri Botanical Garden

M. W. ALEXANDER,
SOUTH-EAST CORNER OF FOURTH AND MARKET STREETS,
ST. LOUIS.

singular structure, inner coating membrane of fruit coating the ovarian cavity, hard, crustaceous, in the ripe fruit breaking to pieces

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

Ribes nigrum both Cultiv April 1860

Ribes rubrum

755



about × 4



oblong lobes of calyx a little longer than the spatulate emarginate or denticulate petals a little shorter than tube, twice as long as ovary; stamens as long as petals style smooth, entire stigma large, bilobed.

orbicular obovate lobes of calyx very much longer than the very short open tube (pateriform) and than the minute petals & short stamens; and ovary stamens longer than petals; styles united to the middle divaricate smooth

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved
MISSOURI BOTANICAL GARDEN

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

4

Ribes

Apl 1860





756

Ribes Grossularia

Cult

English gooseberry

ovate oblong laciniae
of calyx longer than
ventricose or globose tube
longer than ovary, the filaments
& styles.
spatulate obtuse petals 1/3 of
lacinia of calyx
stamens 2/3 of laciniae
styles as long or usually a little
longer, divided almost to base

dr)

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

Ribes floridum L'Her. June 1862

X4

757



Fort Pierre





tube naked inside
lobes about equal to tube
somewhat longer than petals
stamens equal to petals
style bifid at tip

perfectly equal to R. nigrum



St Louis

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Ribes Cereum (Fendler) Jun 1862

× 4

758





Tube perfectly smooth inside
petals inserted below the throat
obovate cucullate, ^s
stamens short, inserted below petals
style naked, divided only at tip
stigma large globose

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

Ribes irriguum Dougl.
Fendler

June 1862

×4







scarcely diffd fr R. hirtellum

759

calyx tube villous at throat only

hairy lobes of calyx longer than funnel-shaped tube, much longer than the smooth ovary (twice as long)
broadly obovate spatulat petals nearly half as long as lobes
stam & style a little shorter than lobes
style divided beyond the middle very hairy part

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI
BOTANICAL
GARDEN

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building Cor. Fourth & Olive Sts., St. Louis, Mo.

Ribes rotundifolia

760

(Hale)

tube of Calyx pubescent inside

lobes of Calyx linear-oblong

twice as long as tube or

as long as tube & ovary together

petals retuse spatulate half

as long as lobes

filaments & style exserted.

(twice as long as lobes of calyx

style hairy below, less than 1/2

divided, at length longer than

stamens.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN